

The Galactolipid GOPO Mediates Beneficial Effects of Rose Hip on Arthritis by Reducing Chemokine and Interleukin Production in Macrophages and Chondrocytes

Joseph Schwager, PhD

Principal Scientist, DSM Nutritional Products

Basel Switzerland

DSM Nutritional Products

APLAR 2008 /JS

Unlimited. DSM


Features:

- Osteoarthritis is a disease condition that affects cartilage (erosion or insufficient regeneration of cartilage).
- Cartilage erosion and rebuilding are highly complex biological processes.
- Rose hip has been found to have beneficial effects in the treatment of OA.
- A constituent galactolipid of rose hip (*i.e.* GOPO) has effects on chemotaxis of neutrophils and thus appears to be one bioactive constituent of rose hip.

Rationale and Experimental Approach:

- Identify bio-actives that account for biological efficacy of rose hip
- Expand biological profiling of one known constituent of rose hip, GOPO
- Establish cellular mode of action of rose hip and GOPO

Structure of GOPO



DSM Nutritional Products

APLAR 2008 /JS

Unlimited. DSM

Cellular models

- Mouse macrophages
 - Permits to determine anti-inflammatory effects
- Peripheral blood leukocytes
 - Permits to measure anti-inflammatory effects on various cell populations
- Chondrocytes (normal human articular cells)
 - Is the most appropriate target cell population to delineate anabolic and catabolic events related to osteoarthritis

Levels of action of rose hip and its constituents

- Inflammatory mediators (PGE₂, nitric oxide; interleukins, chemokines, cytokines)
- Expression of genes (inflammatory genes in macrophages, leukocytes; catabolic and anabolic genes in chondrocytes)




Macrophages

- Inflammatory mediators
 - Nitric oxide
 - PGE₂: pro-inflammatory prostaglandin

Compound	IC ₅₀ (mg/mL)	
	NO	PGE ₂
RH powder	833 ± 40	541 ± 41
GOPO	28 ± 5	>38

Expression of genes (inflammatory genes, chemokines)

- GOPO and RH differentially modulate the expression of iNOS (→ nitric oxide), CXCL-10/IP-10, und CCL5 (→RANTES). The chemokines regulate migration of T_H lymphocytes and macrophages



Approach

- Generate culture supernatants of activated macrophages
- Determine interleukins, chemokines and cytokines by multiparametric ELISA in supernatants

Main features (in mouse macrophages)


- Substantial reduction of the secretion of pro-inflammatory interleukins (IL-1) and concomitant upregulation of anti-inflammatory IL-10
- Only marginal impact on murine chemokines
- No effects on murine TNF- α and INF- γ

		GOPO	RHP
Chemokines	CCL5/RANTES	→	→
	MIP-1alpha	→	→
	MIP-1beta	→	→
	Eotaxin	→	↓
	MCP-1	→	↑
Interleukins	IL-1alpha	↓	↓
	IL-1beta	↓	↓
	IL-4	→	→
	IL-6	→	↓
	IL-10	→	↑
	IL-12	→	→
	TNF-alpha	→	→
IFN-gamma	→	→	

Decreasing: ↓ increasing: ↑ unaltered: →

Approach

- Generate culture supernatants of activated human peripheral blood leukocytes
- Determine interleukins/chemokines/cytokines by multiparametric ELISA



Main features

- The production of key chemokines involved in mobilisation of T_H lymphocytes are reduced by GOPO and/or rose hip
- The production of a key pro-inflammatory interleukin, IL-6 is decreased by GOPO and RH

		GOPO	RHP
Chemokines	CXC10/IP-10	↓	↓
	CCL5/RANTES	→	↓
	MIP-1alpha	↓	→
	MIP-1beta	↓	→
	IL-8	↓	→
	Eotaxin	↓	→
	MCP-1	↓	↑
Interleukins	IL-1beta	↓	↓
	IL4	↓	→
	IL-6	↓	↓
	IL-10	↓	→
	IL-12	↓	↓
	TNF-alpha	↓	↓
	IFN-gamma	↓	↓

Main features


- Various chemokines are robustly reduced by GOPO and to a lower extent by rose hip powder
- Both substances have inhibitory effects on a panel of pro-inflammatory interleukins and cytokines

Decreasing: ↓ increasing: ↑ unaltered: →


DSM Nutritional Products

Approach

- Stimulate human PBL in the presence of graded amounts of GOPO or RH
- Determine expression of genes by quantitative RT-PCR



LPS/IFN-g-only: 277-fold







LPS/IFN-g-only: 35960-fold

Main features

- GOPO differentially modulate the expression of the pro-inflammatory interleukin IL1- β and IL-6.
- The effect of rose hip on the expression of these genes is much weaker.

Approach

- NHAC-kn were activated with IL-1 β (representing ,diseased‘ joints) in the presence of RHP or GOPO for 24 hours. The chemokine or PGE₂ production was determined by multtiparametric RT-PCR




Main features

- GOPO dose-dependently inhibits the synthesis of three chemokines
- GOPO dosedependently inhibits the production of the pro-inflammatory PGE₂

Approach

- NHAC-kn were activated with IL-1 β (representing ,diseased‘ joints) in the presence of RHP or GOPO for 4 hours. The gene expression was measured by quantitative RT-PCR.



Main features

- Expression of aggrecanase and matrix metalloproteinase that trigger erosion of joints are dose-dependently reduced by GOPO and RH


DSM Nutritional Products

APLAR 2008 /JS

Unlimited. DSM

Experimental approach

- NHAC-kn were cultured in the presence of RHP (50 mg/L) or GOPO (19 mg/L) for 4 hours. The gene expression was measured by quantitative RT-PCR (in comparison to unstimulated chondrocytes, representing „healthy“ joints).



Main features

- Anabolic genes that contribute to the expression of the extracellular matrix proteins (e.g. aggrecan and collagen) are activated by GOPO and RH
- Similarly, a transcription factor (SOX-9) that is involved in chondrocyte proliferation and differentiation is up-regulated by the compounds

- Bio-actives have been identified in rose hip powder by monitoring anti-inflammatory parameters and multi-parametric profiling in three cellular systems.
- Mouse macrophages were convenient for identifying first effects, while the heterogenous PBL populations (containing mononuclear and polymorphonuclear cells) permitted to monitor more diverse *i.e.* pleiotropic effects.
- GOPO has been identified as a potent effector molecule that modulates various facets of the inflammatory processes and cell migration mediated by chemokines.
- The importance of GOPO in cartilage protection is suggested by its effect on the chemokine production by chondrocytes and the expression of catabolic and anabolic genes by human articular cells.
- Although RH contains significant quantities of GOPO (<0.1%), the contents cannot account for the whole biological activity of RH.
- Consequently, other constituents contribute to, and might act in concert, to reduce the erosion of the extracellular matrix in joints or favor the rebuilding of cartilage.